arXiv:0801.2207v1 [math.RA] 15 Jan 2008

The derivation algebra and automorphism group of the twisted Schrödinger-Virasoro algebra[1]

Junbo Li*,†), Yucai Su‡)

*)Department of Mathematics, Shanghai Jiao Tong University, Shanghai 200240, China

†)Department of Mathematics, Changshu Institute of Technology, Changshu 215500, China

‡)Department of Mathematics, University of Science and Technology of China, Hefei 230026, China

E-mail: sd_junbo@163.com, ycsu@ustc.edu.cn

Abstract. In this article, we determine the derivation algebra Der$\mathscr{L}$ and the automorphism group Aut$\mathscr{L}$ of the twisted Schrödinger-Virasoro algebra $\mathscr{L}$.

Key words: twisted Schrödinger-Virasoro algebra; derivation algebra; automorphism group.

MR(2000) Subject Classification: 17B05, 17B40, 17B65, 17B68.

1. Introduction

It is well known that the infinite-dimensional Schrödinger Lie algebras and Virasoro algebra play important roles in many areas of mathematics and physics (e.g., statistical physics). The original Schrödinger-Virasoro Lie algebra was introduced in [5], in the context of non-equilibrium statistical physics, containing as subalgebras both the Lie algebra of invariance of the free Schrödinger equation and the Virasoro algebra. The infinite-dimensional Lie algebra discussed in this article called the twisted Schrödinger-Virasoro algebra is the twisted deformation of the original Schrödinger-Virasoro Lie algebra.

Both original and twisted Schrödinger-Virasoro Lie algebras are closely related to the Schrödinger Lie algebras and the Virasoro Lie algebra (see [6], [7], [14] and [12]). They should consequently play a role akin to that of the Virasoro Lie algebra in two-dimensional equilibrium statistical physics.

Motivated by the research for deformations and central extensions of both original and twisted Schrödinger-Virasoro Lie algebras, the sets of generators provided by the cohomology classes of the cocycles were presented in [10]. In [13], the author constructed vertex algebra representations of the Schrödinger-Virasoro Lie algebras out of a charged symplectic boson and a free boson with its associated vertex operators. The purpose of this article is to determine the derivation algebra and the automorphism group of the twisted Schrödinger-Virasoro Lie algebras $\mathscr{L}$. We show that the derivation algebra of $\mathscr{L}$ is the direct sum of three linear independent outer derivations and the inner derivation algebra. Finally, we characterize the automorphism group Aut$\mathscr{L}$ of $\mathscr{L}$.

Now we give the definition of the Lie algebra $\mathscr{L}$. A Lie algebra $\mathscr{L}$ is called a *twisted*

---

[1] Supported by NSF grants 10471091, 10671027 of China, One Hundred Talents Program from University of Science and Technology of China.

Schrödinger-Virasoro Lie algebra (see [10]), if $\mathscr{L}$ has the $\mathbb{C}$-basis

$$\{L_n, Y_n, M_n, C \,|\, n \in \mathbb{Z}\}$$

with the Lie brackets

$$[L_n, L_{n'}] = (n' - n)L_{n+n'} + \delta_{n,-n'}\frac{n^3 - n}{12}C, \tag{1.1}$$

$$[L_n, Y_m] = (m - \frac{n}{2})Y_{n+m}, \tag{1.2}$$

$$[L_n, M_p] = pM_{n+p}, \tag{1.3}$$

$$[Y_m, Y_{m'}] = (m' - m)M_{m+m'}, \tag{1.4}$$

$$[Y_m, M_p] = [M_n, M_p] = [\mathscr{L}, C] = 0. \tag{1.5}$$

The twisted Schrödinger-Virasoro Lie algebra has an infinite-dimensional *twisted Schrödinger subalgebra* denoted by SS with the $\mathbb{C}$-basis $\{Y_n, M_n \,|\, n \in \mathbb{Z}\}$ and a *Virasoro subalgebra* denoted by $\mathfrak{V}$ with the $\mathbb{C}$-basis $\{L_n, C \,|\, n \in \mathbb{Z}\}$. The action of the Virasoro subalgebra on the Schrödinger subalgebra is natural. The center of $\mathscr{L}$, denoted by $C(\mathscr{L})$, is two-dimensional, spanned by $\{M_0, C\}$. Introduce a $\mathbb{Z}$-gradation on $\mathscr{L}$ by $\mathrm{deg}L_n = \mathrm{deg}Y_n = \mathrm{deg}M_n = n$, $\mathrm{deg}C = 0$ and decompose $\mathscr{L}$ with respect to the following gradation:

$$\mathscr{L} = \bigoplus_{n\in\mathbb{Z}} \mathscr{L}_n,\ \mathscr{L}_0 = \mathrm{Span}_{\mathbb{C}}\{L_0, Y_0, M_0, C\},\ \mathscr{L}_n = \mathrm{Span}_{\mathbb{C}}\{L_n, Y_n, M_n\},\ n \in \mathbb{Z}\backslash\{0\}.$$

Throughout the article, we denote by $\mathbb{Z}^*$ the set of all nonzero integers and $\mathbb{C}^*$ the set of all nonzero complex numbers.

2. The derivation algebra Der$\mathscr{L}$ of $\mathscr{L}$

Let V be a $\mathscr{L}$-module. A linear map φ from $\mathscr{L}$ to V is called a *derivation*, if for any $x, y \in \mathscr{L}$, we have

$$\varphi[x, y] = x.\varphi(y) - y.\varphi(x).$$

For $v \in V$, the map $\phi : x \to x.v$ is called an *inner derivation*.

Denote by $\mathrm{Der}(\mathscr{L}, V)$ the vector space of all derivations, $\mathrm{Inn}(\mathscr{L}, V)$ the vector space of all inner derivations. Then the first cohomology group of $\mathscr{L}$ with coefficients in V is

$$H^1(\mathscr{L}, V) = \mathrm{Der}_{\mathbb{C}}(\mathscr{L}, V)/\mathrm{Inn}_{\mathbb{C}}(\mathscr{L}, V). \tag{2.1}$$

The right-hand side is also called the space of *outer derivations*.

By definition, the algebra $\mathscr{L}$ is a semidirect product of $\mathbb{Z}$-graded algebras $\mathscr{L} = \mathfrak{V} \ltimes SS$, with $\mathfrak{V} := \bigoplus_{n\in\mathbb{Z}} \mathbb{C}L_n \oplus \mathbb{C}C$ being the Virasoro algebra and $SS := \bigoplus_{n\in\mathbb{Z}}(\mathbb{C}Y_n \oplus \mathbb{C}M_n)$ being the twisted Schrödinger algebra. Clearly, $\mathfrak{V}$ is $\mathbb{Z}$-graded by $\mathfrak{V}_0 = \mathbb{C}L_0 \oplus \mathbb{C}C$ and $\mathfrak{V}_n = \mathbb{C}L_n$ for $n \in \mathbb{Z} \setminus \{0\}$ while SS is $\mathbb{Z}$-graded by $SS_n = \mathbb{C}Y_n \oplus \mathbb{C}M_n$ for $n \in \mathbb{Z}$. By [8], the short exact sequence

$$\{0\} \longrightarrow SS \longrightarrow \mathscr{L} \longrightarrow \mathscr{L}/SS \longrightarrow \{0\},$$

induces a long exact sequence

$$\begin{aligned}\{0\} \longrightarrow \quad & H^0(\mathscr{L}, SS) \longrightarrow H^0(\mathscr{L}, \mathscr{L}) \xrightarrow{f} H^0(\mathscr{L}, \mathscr{L}/SS) \longrightarrow \\ & H^1(\mathscr{L}, SS) \longrightarrow H^1(\mathscr{L}, \mathscr{L}) \longrightarrow H^1(\mathscr{L}, \mathscr{L}/SS) \longrightarrow \cdots \end{aligned} \tag{2.2}$$

of $\mathbb{Z}$-graded vector spaces. Note that $H^0(\mathscr{L}, SS) = SS^{\mathscr{L}} = \mathbb{C}M_0$, while $H^0(\mathscr{L}, \mathscr{L}) = \mathscr{L}^{\mathscr{L}} \cong \mathbb{C}M_0 + \mathbb{C}C$, and $H^0(\mathscr{L}, \mathscr{L}/SS) = (\mathscr{L}/SS)^{\mathscr{L}} \cong \mathbb{C}C$. From this we see that the map f is surjective. Thus (2.2) gives the following exact sequence

$$\{0\} \longrightarrow H^1(\mathscr{L}, SS) \longrightarrow H^1(\mathscr{L}, \mathscr{L}) \longrightarrow H^1(\mathscr{L}, \mathscr{L}/SS). \tag{2.3}$$

The right-hand side of (2.3) can be computed from the exact sequence

$$\{0\} \longrightarrow H^1(\mathscr{L}/SS, \mathscr{L}/SS) \longrightarrow H^1(\mathscr{L}, \mathscr{L}/SS) \longrightarrow H^1(SS, \mathscr{L}/SS)^{\mathscr{L}}. \tag{2.4}$$

Note that $H^1(\mathscr{L}/SS, \mathscr{L}/SS) = H^1(\mathfrak{V}, \mathfrak{V})$ is the algebra of outer derivations of the Virasoro algebra $\mathfrak{V}$, which is equal to zero by [14], while $H^1(SS, \mathscr{L}/SS)^{\mathscr{L}}$ can be embeded into $\mathrm{Hom}_{\mathcal{U}(\mathfrak{V})}(SS/[SS, SS], \mathfrak{V})$ (where $\mathcal{U}(\mathfrak{V})$ is the universal enveloping algebra of $\mathfrak{V}$). Therefore, by (2.3) and (2.4), $H^1(\mathscr{L}, \mathscr{L})$ will follow from the result of the computation of $H^1(\mathscr{L}, SS)$ and $\mathrm{Hom}_{\mathcal{U}(\mathfrak{V})}(SS/[SS, SS], \mathfrak{V})$.

Lemma 2.1. $\mathrm{Der}(\mathscr{L}, SS) = \mathrm{Der}(\mathscr{L}, SS)_0 + \mathrm{Inn}(\mathscr{L}, SS)$, *where*

$$\mathrm{Der}(\mathscr{L}, SS)_0 = \{\mathcal{D} \in \mathrm{Der}(\mathscr{L}, SS) \mid \mathcal{D}(\mathscr{L}_n) \subseteq SS_n, \ \textit{for any } n \in \mathbb{Z}\}.$$

Proof. This follows immediately from (1.2) of [1]. □

Lemma 2.2. *$\mathcal{D} \in \mathrm{Der}(\mathscr{L}, SS)_0$ if and only if $\mathcal{D}$ is a linear map from $\mathscr{L}$ to SS satisfying the following conditions:*

$$\begin{aligned} &\text{(i)} && \mathcal{D}(L_n) = \big(dn + d_1\big)M_n, \\ &\text{(ii)} && \mathcal{D}(M_n) = 2g_0 M_n, \\ &\text{(iii)} && \mathcal{D}(Y_n) = g_0 Y_n, \\ &\text{(iv)} && \mathcal{D}(C) = 0, \end{aligned}$$

for any $n \in \mathbb{Z}$ and some d, d_1, $g_0 \in \mathbb{C}$.

Proof. Let $\mathcal{D}$ be a linear map form $\mathscr{L}$ to SS satisfying (i)–(iv). It is clear that $\mathcal{D}(\mathscr{L}_n) \subseteq SS_n$, and it is easy to check that $\mathcal{D}$ is a derivation from $\mathscr{L}$ to SS.

Let $\mathcal{D} \in \mathrm{Der}(\mathscr{L}, SS)_0$. For any $n \in \mathbb{Z}$, assume

$$\mathcal{D}(L_n) = c_n Y_n + d_n M_n, \qquad c_n,\, d_n \in \mathbb{C}. \tag{2.5}$$

Applying $\mathcal{D}$ to $[L_1, L_n] = (n-1)L_{n+1}$ and $[L_{-1}, L_{n+1}] = (n+2)L_n$, we obtain

$$\big(d_1 - nd_n + (n-1)d_{n+1}\big)M_{n+1} = \big((\frac{n}{2} - 1)c_1 + (n - \frac{1}{2})c_n - (n-1)c_{n+1}\big)Y_{n+1}, \tag{2.6}$$

$$\big((n+2)d_n - d_{-1} - (n+1)d_{n+1}\big)M_n = \big(\frac{n+3}{2}c_{-1} - (n+2)c_n + (n + \frac{3}{2})c_{n+1}\big)Y_n. \tag{2.7}$$

Comparing the coefficients of M_{n+1}, Y_{n+1} in (2.6) and M_n, Y_n in (2.7), one has

$$(n-1)d_{n+1} = nd_n - d_1, \tag{2.8}$$

$$(n-1)c_{n+1} = (\frac{n}{2} - 1)c_1 + (n - \frac{1}{2})c_n, \tag{2.9}$$

$$(n+1)d_{n+1} = (n+2)d_n - d_{-1}, \tag{2.10}$$

$$(n + \frac{3}{2})c_{n+1} = (n+2)c_n - \frac{n+3}{2}c_{-1}. \tag{2.11}$$

Subtracting (2.8) from (2.10), and (2.9) from (2.11), we respectively obtain

$$d_{n+1} - d_n = \frac{d_1 - d_{-1}}{2}, \tag{2.12}$$

$$c_{n+1} - c_n = -\frac{n}{5}(c_{-1} + c_1) + \frac{1}{5}(2c_1 - 3c_{-1}). \tag{2.13}$$

Denoting $\frac{d_1 - d_{-1}}{2} = d$ and using induction on n in (2.12), one can deduce

$$d_n = d_1 + (n-1)d, \;\; \forall\, n \in \mathbb{Z}. \tag{2.14}$$

Taking $n = -1$ in (2.11), one obtains $c_0 = 0$. Then taking $n = 0$ in (2.13) gives $c_1 = -c_{-1}$. From this, (2.13) can be rewritten as $c_{n+1} - c_n = c_1$. Thus $c_n = nc_1$. This together with (2.9) gives

$$c_n = 0, \quad \forall\, n \in \mathbb{Z}. \tag{2.15}$$

According to the equations (2.14) and (2.15), for any $n \in \mathbb{Z}$, we can rewrite (2.5) as

$$\mathcal{D}(L_n) = \big(d_1 + (n-1)d\big)M_n, \;\; \text{where} \;\; d,\, d_1 \in \mathbb{C}. \tag{2.16}$$

Re-denoting $d_1 - d$ by d_1, we obtain (i).

Applying $\mathcal{D}$ to $[L_{-3}, L_3] = 6L_0 - 2C$, one has

$$\begin{aligned}
6(d_1 - d)M_0 - 2\mathcal{D}(C) &= 6\mathcal{D}(L_0) - 2\mathcal{D}(C) = \mathcal{D}\big([L_{-3}, L_3]\big) \\
&= [(d_1 - 4d)M_{-3}, L_3] + [L_{-3}, (d_1 + 2d)M_3] \\
&= 3(d_1 - 4d)M_0 + 3(d_1 + 2d)M_0 \\
&= 6(d_1 - d)M_0.
\end{aligned}$$

This gives $\mathcal{D}(C) = 0$. Thus (iv) follows.

For any $n \in \mathbb{Z}$, write

$$\mathcal{D}(Y_n) = g_n Y_n + h_n M_n, \quad g_n,\, h_n \in \mathbb{C}. \tag{2.17}$$

Applying $\mathcal{D}$ to $[L_1, Y_n] = (n - \frac{1}{2})Y_{n+1}$ and comparing the coefficients of Y_{n+1} and M_{n+1}, we obtain

$$(n - \frac{1}{2})g_n = (n - \frac{1}{2})g_{n+1}, \quad nh_n = (n - \frac{1}{2})h_{n+1}. \tag{2.18}$$

Applying $\mathcal{D}$ to $[L_{-1}, Y_1] = \frac{3}{2}Y_0$ and comparing the coefficients of Y_0 and M_0, we obtain $g_1 = g_0$, $h_1 = \frac{3}{2}h_0$. Then using induction on n in (2.18), one obtains

$$g_n = g_0, \quad h_n = 0, \ \ \forall\ n \in \mathbb{Z}. \tag{2.19}$$

Therefore, for any $n \in \mathbb{Z}$, one can rewrite (2.17) as $\mathcal{D}(Y_n) = g_0 Y_n$, i.e., (iii) follows.

For any $m, n \in \mathbb{Z}$, applying $\mathcal{D}$ to $[Y_m, Y_n] = (n - m)M_n$ and using (iii), one can easily obtain (ii). $\square$

Set $d_1 = 1$, $d = g_0 = 0$; $d = 1$, $d_1 = g_0 = 0$; $g_0 = 1$, $d_1 = d = 0$ in $\mathcal{D} \in \mathrm{Der}(\mathscr{L}, SS)_0$ respectively, we obtain three derivations from $\mathscr{L}$ to SS:

$$\begin{aligned}
&\mathcal{D}_1: \ \mathcal{D}_1(L_n) = M_n, && \mathcal{D}_1(M_n) = \mathcal{D}_1(Y_n) = \mathcal{D}_1(C) = 0; \\
&\mathcal{D}_2: \ \mathcal{D}_2(L_n) = nM_n, && \mathcal{D}_2(M_n) = \mathcal{D}_2(Y_n) = \mathcal{D}_2(C) = 0; \\
&\mathcal{D}_3: \ \mathcal{D}_3(Y_n) = Y_n, && \mathcal{D}_3(M_n) = 2M_n,\ \mathcal{D}_3(L_n) = \mathcal{D}_3(C) = 0;
\end{aligned}$$

where $n \in \mathbb{Z}$. It is straightforward to verify that $\mathcal{D}_i$, $i = 1, 2, 3$, are linear independent outer derivations. Thus (2.1) and Lemma 2.2 prove Lemma 2.3 below.

Lemma 2.3. $H^1(\mathscr{L}, SS) = \mathbb{C}\mathcal{D}_1 \dot{+} \mathbb{C}\mathcal{D}_2 \dot{+} \mathbb{C}\mathcal{D}_3$.

Lemma 2.4. $\mathrm{Hom}_{\mathcal{U}(\mathfrak{V})}(SS/[SS, SS], \mathfrak{V}) = 0$.

Proof. By the definition of SS, we have $[SS, SS] = \mathrm{Span}_{\mathbb{C}}\{M_n \,|\, n \in \mathbb{Z}\}$. For any $f \in \mathrm{Hom}_{\mathcal{U}(\mathfrak{V})}(SS/[SS, SS], \mathfrak{V})$ and any $n \in \mathbb{Z}$, one can assume

$$f(Y_n) = \sum_k p_k^{(n)} L_k + c^{(n)} C, \quad \text{for some } \ p_k^{(n)}, \ c^{(n)} \in \mathbb{C}.$$

For any $m, n \in \mathbb{Z}$, applying f to both sides of $[L_m, Y_n] = (n-m)Y_{m+n}$, using $f([L_m, Y_n]) = [L_m, f(Y_n)]$, one has

$$\sum_k (k-m) p_k^{(n)} L_{k+m} + p_{-m}^{(n)} \frac{m^3 - m}{12} C = (n-m)\big(\sum_k p_k^{(m+n)} L_k + c^{(m+n)} C\big). \tag{2.20}$$

Comparing the coefficients of C and L_k in (2.20), one has

$$p_{-m}^{(n)} \frac{m^3 - m}{12} = (n-m) c^{(m+n)}, \tag{2.21}$$

$$(n-m) p_k^{(m+n)} = (k - 2m) p_{k-m}^{(n)}. \tag{2.22}$$

Taking $m = 0$, $n \neq 0$ and $m = 1, n = -1$ in (2.21) respectively, one immediately obtains $C^{(n)} = 0$ for $n \in \mathbb{Z}$. Using this in (2.21), one has

$$p_k^{(n)} = 0 \quad \text{for all} \quad k \notin \{-1, 0, 1\}. \tag{2.23}$$

If $k \in \{-1, 0, 1\}$, by choosing some m such that $2n - m \neq 0$ and $k - m \notin \{-1, 0, 1\}$ and replacing n by $n - m$ in (2.22), and using (2.23), one again obtains $p_k^{(n)} = 0$. This proves $f(Y_n) = 0$ for all $n \in \mathbb{Z}$. □

Theorem 2.5. *$Der\mathscr{L} = \mathbb{C}\mathcal{D}_1 \dotplus \mathbb{C}\mathcal{D}_2 \dotplus \mathbb{C}\mathcal{D}_3 \dotplus ad\mathscr{L}$.*

Proof. By Lemma 2.4, $H^1(SS, \mathscr{L}/SS)^{\mathscr{L}} = 0$. According to the exact sequence (2.4) and the fact $H^1(\mathscr{L}/SS, \mathscr{L}/SS) = 0$, we have $H^1(\mathscr{L}, \mathscr{L}/SS) = 0$. So by Lemma 2.3 and the exact sequence (2.3), one has

$$H^1(\mathscr{L}, \mathscr{L}) = \mathbb{C}\mathcal{D}_1 \dotplus \mathbb{C}\mathcal{D}_2 \dotplus \mathbb{C}\mathcal{D}_3.$$

Hence the theorem follows from $H^1(\mathscr{L}, \mathscr{L}) = \mathrm{Der}\mathscr{L}/\mathrm{ad}\mathscr{L}$. □

3. The automorphism group $\mathrm{Aut}\mathscr{L}$ of $\mathscr{L}$

Throughout this section, we denote by $\mathrm{Inn}(\mathscr{L})$ the set of inner automorphisms. Then $\mathrm{Inn}(\mathscr{L})$ is a normal subgroup of $\mathrm{Aut}\mathscr{L}$ and

$$\mathrm{Inn}(\mathscr{L}) \text{ is generated by } \exp(k\mathrm{ad}L_0), \ \exp(k\mathrm{ad}Y_n) \text{ and } \exp(k\mathrm{ad}M_n), \tag{3.1}$$

where $n \in \mathbb{Z}, k \in \mathbb{C}$.

Denote $J_1 = \mathrm{Span}_{\mathbb{C}}\{M_n, C \,|\, n \in \mathbb{Z}\}$ and $J_2 = \mathrm{Span}_{\mathbb{C}}\{Y_n, M_n, C \,|\, n \in \mathbb{Z}\}$. They both are ideals of $\mathscr{L}$. And the following lemma is clear.

Lemma 3.1. *For any* $ß \in Aut\mathscr{L}$*, one has* $ß(J_i) \subseteq J_i\,(i=1,2)$ *and* $ß\big(C(\mathscr{L})\big) \subseteq C(\mathscr{L})$*.*

For any $a \in \mathbb{C}^*$, we can define the following maps on $\mathscr{L}$.

$$\begin{aligned}
ß_a:&\quad L_n \longrightarrow a^n L_n, \quad Y_n \longrightarrow a^n Y_n, \quad M_n \longrightarrow a^n M_n, \quad C \longrightarrow C; && (3.2)\\
\varepsilon:&\quad L_n \longrightarrow -L_{-n}, \quad Y_n \longrightarrow -Y_{-n}, \quad M_n \longrightarrow -M_{-n}, \quad C \longrightarrow -C; && (3.3)\\
\psi_a:&\quad L_n \longrightarrow L_n, \quad Y_n \longrightarrow aY_n, \quad M_n \longrightarrow a^2 M_n, \quad C \longrightarrow C. && (3.4)
\end{aligned}$$

It is easy to see that $ß_a \in \mathrm{Inn}\mathscr{L},\ \varepsilon, \psi_a \in \mathrm{Aut}\mathscr{L} \setminus \mathrm{Inn}\mathscr{L}$.

Lemma 3.2. *For any* $ß \in \mathrm{Aut}(\mathscr{L})$*, there exists an automorphism* $\xi \in \mathrm{Inn}(\mathscr{L})$ *of the form* $\exp\big(\sum_{j\neq 0} b'_j \mathrm{ad}Y_j + \sum_{k\neq 0} c'_k \mathrm{ad}M_k\big)$ *such that* $\xi^{-1}\sigma(L_0) = a(L_0 + b_0Y_0 + c_0M_0 + d_0C)$*, where* $a \in \mathbb{C}^*, b_0, c_0, d_0 \in \mathbb{C}$*.*

Proof. For any $ß \in \mathrm{Aut}(\mathscr{L})$, we can assume that

$$\sigma(L_0) = \sum_{i\neq 0} a_i L_i + \sum_{j\neq 0} b_j Y_j + \sum_{k\neq 0} c_k M_k + a(L_0 + b_0Y_0 + c_0M_0 + d_0C), \tag{3.5}$$

where $a_i, b_j, c_k \in \mathbb{C}$, $a \in \mathbb{C}^*$ (we must have $a \neq 0$, otherwise, $\sigma(L_0)$ would be ad-locally nilpotent while L_0 is ad-semi-simple).

We shall find some $\xi = \exp\big(\sum_{j\neq 0} b'_j \mathrm{ad}Y_j + \sum_{k\neq 0} c'_k \mathrm{ad}M_k\big) \in \mathrm{Inn}(\mathscr{L})$, such that

$$\sigma(L_0) = \exp\big(\sum_{j\neq 0} b'_j \mathrm{ad}Y_j + \sum_{k\neq 0} c'_k \mathrm{ad}M_k\big)\big(a'(L_0 + b'_0Y_0 + c'_0M_0 + d'_0C)\big). \tag{3.6}$$

for some $a' \in \mathbb{C}^*, b'_j, c'_k, d'_0 \in \mathbb{C}$. We only need to solve a', b'_j, c'_k, d'_0 by comparing the coefficients of L_i, Y_j, M_k and C, which gives

$$a = a',\ b_0 = b'_0,\ c_0 = c'_0,\ d_0 = d'_0,\ b'_j = -a^{-1}j^{-1}b_j, \tag{3.7}$$

$$c'_k = -b_0b'_k - a^{-1}k^{-1}c_k - \frac{\sum\limits_{i+l=k} l(l-i)k^{-1}a^{-2}i^{-1}l^{-1}b_ib_l}{2}. \tag{3.8}$$

Applying ξ^{-1} to both sides of (3.5) and using (3.7), we obtain

$$\xi^{-1}\sigma(L_0) = a(L_0 + b_0Y_0 + c_0M_0 + d_0C). \tag{3.9}$$

Lemma 3.3. $b_0 = 0$.

Proof. Since L_0 is an ad-semi-simple element in $\mathscr{L}$, then $\xi^{-1}\sigma(L_0) = a(L_0 + b_0Y_0 + c_0M_0 + d_0C)$ is also ad-semi-simple in $\mathscr{L}$. By linear algebra, the following matrix (given by the adjoint action of $\operatorname{ad} a(L_0 + b_0Y_0 + c_0M_0 + d_0C)$ on the basis $\{L_n, Y_n, M_n, C\}$ of $\mathscr{L}_n$)

$$\begin{pmatrix} an & 0 & 0 & 0 \\ \frac{ab_0n}{2} & an & 0 & 0 \\ 0 & ab_0n & an & 0 \\ 0 & 0 & 0 & 0 \end{pmatrix}$$

can be diagonalized, which forces $b_0 = 0$. This lemma follows. $\square$

According to Lemma 3.2 and 3.3, for any $\text{ß} \in \operatorname{Aut}(\mathscr{L})$, by replacing ß with $\xi\text{ß}$ for some $\xi \in \operatorname{Inn}(\mathscr{L})$, we can write

$$\text{ß}(L_0) = a(L_0 + c_0M_0 + d_0C), \tag{3.10}$$

where $a \in \mathbb{C}^*, c_0, d_0 \in \mathbb{C}$.

Lemma 3.4. *For any $\text{ß} \in \operatorname{Aut}(\mathscr{L})$, by replacing ß with $\varepsilon\text{ß}$ if necessary where ε is defined in (3.3), we can write*

$$\text{ß}(L_0) = L_0 + c_0M_0 + d_0C,\ \text{ß}(Y_n) = e'_nY_n + e''_nM_n + \delta_{n,0}e_0C,\ \text{ß}(M_n) = f'_nM_n + \delta_{n,0}f_0C,$$

for some $n \in \mathbb{Z}$, e'_n, $f'_n \in \mathbb{C}^*$, $e_0, f_0, e''_n \in \mathbb{C}$.

Proof. By Lemma 3.1, we can write

$$\text{ß}(Y_n) = \sum_i e'_iY_i + \sum_j e''_jM_j + e_nC, \tag{3.11}$$

$$\text{ß}(M_n) = \sum_i f'_iM_i + f_nC, \tag{3.12}$$

where $e'_i \in \mathbb{C}^*, f'_i \in \mathbb{C}^*\ (i \neq 0)$, $e''_j, e_n, f'_0, f_n \in \mathbb{C}$, $n \in \mathbb{Z}$.

Applying ß to $[L_0, Y_n] = nY_n$ and $[L_0, M_n] = nM_n$, we obtain

$$\Big[a(L_0 + c_0M_0 + d_0C), \sum_i e'_iY_i + \sum_j e''_jM_j + e_nC\Big] = n\big(\sum_i e'_iY_i + \sum_j e''_jM_j + e_nC\big),$$

$$\Big[a(L_0 + c_0M_0 + d_0C), \sum_i f'_iM_i + f_nC\Big] = n\big(\sum_i f'_iM_i + f_nC\big).$$

That is,

$$\sum_i (ai - n)e'_iY_i + \sum_j (aj - n)e''_jM_j - ne_nC = 0, \tag{3.13}$$

$$\sum_i (ai - n)f'_iM_i - nf_nC = 0. \tag{3.14}$$

Comparing the coefficients of Y_i, M_j and C in the above two equations, we obtain

$$i = \frac{n}{a} \in \mathbb{Z}, \ \ (aj-n)e''_j = 0, \ \ e_n = \delta_{n,0}e_0 \ \text{ and } \ f_n = \delta_{n,0}f_0, \tag{3.15}$$

for all $n \in \mathbb{Z}$. It follows that

$$a = \pm 1, \ \ i = an \ \text{ and } \ e''_j = 0 \ \text{ if } \ j \neq an. \tag{3.16}$$

According to (3.15) and (3.16), one can respectively rewrite (3.11) and (3.12) as

$$\text{ß}(Y_n) = e'_{an}Y_{an} + e''_{an}M_{an} + \delta_{n,0}e_0C, \tag{3.17}$$

$$\text{ß}(M_n) = f'_{an}M_{an} + \delta_{n,0}f_0C. \tag{3.18}$$

For one case $a = 1$, the lemma is right. For the other case $a = -1$, replacing ß with εß, we can rewrite (3.10), (3.17) and (3.18) as those in the lemma. Then this lemma follows. □

Lemma 3.5. *Let* ß $\in \mathrm{Aut}(\mathscr{L})$ *be such that*

$$\text{ß}(L_0) = L_0 + c_0M_0 + d_0C, \ \text{ß}(Y_n) = e'_nY_n + e''_nM_n + \delta_{n,0}e_0C, \ \text{ß}(M_n) = f'_nM_n + \delta_{n,0}f_0C,$$

where $n \in \mathbb{Z}$, e'_n, $f'_n \in \mathbb{C}^*$, $e_0, f_0, e''_n \in \mathbb{C}$. *Then by replacing* ß *by* ψ_xß$_y$ß *for some* $x, y \in \mathbb{Z}^*$ (*where* ß$_y$ *and* ψ_x *are defined in* (3.2) *and* (3.4) *respectively*), *one can suppose* ß $= \delta_{\alpha,\beta,\gamma}$ *for some* $\alpha, \beta, \gamma \in \mathbb{C}$, *where* $\delta_{\alpha,\beta,\gamma}$ *is defined by*

$$\delta_{\alpha,\beta,\gamma}(C) = C, \tag{3.19}$$

$$\delta_{\alpha,\beta,\gamma}(M_n) = M_n, \tag{3.20}$$

$$\delta_{\alpha,\beta,\gamma}(Y_n) = Y_n + 2\alpha n M_n, \tag{3.21}$$

$$\delta_{\alpha,\beta,\gamma}(L_n) = L_n + \alpha n Y_n + \big(\alpha^2n^2 + \beta n + \gamma\big)M_n. \tag{3.22}$$

Proof. For any $n \in \mathbb{Z}^*$, we can write

$$\sigma(L_n) = \sum_i u_iL_i + \sum_j v_jY_j + \sum_{k\neq 0} w_kM_k + C_n, \tag{3.23}$$

where $u_i \in \mathbb{C}^*, v_j, w_k \in \mathbb{C}, \ C_n \in C(\mathscr{L})$.

For any $n \in \mathbb{Z}^*$, applying σ to $nL_n = [L_0, L_n]$, we obtain

$$\begin{aligned}\sum_i nu_iL_i + \sum_j nv_jY_j + \sum_{k\neq 0} nw_kM_k + nC_n &= n\sigma(L_n) = \sigma\big([L_0, L_n]\big)\\ &= \sum_i iu_iL_i + \sum_j jv_jY_j + \sum_{k\neq 0} kw_kM_k.\end{aligned}$$

Comparing the coefficients of L_i, Y_i, M_i and C_n, one has

$$u_i = v_i = w_i = 0 \ \text{ if } \ i \neq n, \ \text{ and } \ C_n = 0 \ \text{ for any } \ n \in \mathbb{Z}^*.$$

Hence (3.23) can be rewritten as

$$\sigma(L_n) = u_n L_n + v_n Y_n + w_n M_n \quad \text{where} \quad n \in \mathbb{Z}^*. \tag{3.24}$$

For $n \neq 0, \pm 1$, applying σ to $(n-1)L_{n+1} = [L_1, L_n]$, we obtain

$$\begin{aligned}(n-1)\sigma(L_{n+1}) &= (n-1)(u_{n+1}L_{n+1} + v_{n+1}Y_{n+1} + w_{n+1}M_{n+1}) = [\sigma(L_1), \sigma(L_n)] \\ &= (n-1)u_1u_nL_{n+1} + \big((n-\frac{1}{2})u_1v_n + (\frac{n}{2}-1)v_1u_n\big)Y_{n+1} \\ &\quad +\big(nu_1w_n + (n-1)v_1v_n - w_1u_n\big)M_{n+1}.\end{aligned}$$

Comparing the coefficients of L_{n+1}, Y_{n+1} and M_{n+1}, one has

$$u_{n+1} = u_1u_n, \tag{3.25}$$

$$(n-1)v_{n+1} = (n-\frac{1}{2})u_1v_n + (\frac{n}{2}-1)v_1u_n, \tag{3.26}$$

$$(n-1)w_{n+1} = nu_1w_n + (n-1)v_1v_n - w_1u_n. \tag{3.27}$$

Applying σ to $[L_1, L_{-1}] = -2L_0$, we obtain

$$\begin{aligned}-2\sigma(L_0) &= -2(L_0 + c_0M_0 + d_0C) = \text{ß}\big([L_1, L_{-1}]\big) \\ &= -2u_1u_{-1}L_0 - \frac{3}{2}(u_1v_{-1} + v_1u_{-1})Y_0 - (u_1w_{-1} + 2v_1v_{-1} + w_1u_{-1})M_0.\end{aligned}$$

Comparing the coefficients of L_0, Y_0, M_0 and C, one has

$$u_{-1} = u_1^{-1}, \tag{3.28}$$

$$v_{-1} = -u_1^{-2}v_1, \tag{3.29}$$

$$c_0 = \frac{1}{2}(u_1w_{-1} + 2v_1v_{-1} + w_1u_1^{-1}), \quad d_0 = 0. \tag{3.30}$$

By (3.10), we can rewrite $\text{ß}L_0$ as

$$\text{ß}(L_0) = L_0 + \frac{1}{2}(u_1w_{-1} + 2v_1v_{-1} + w_1u_1^{-1})M_0.$$

Applying σ to $[L_2, L_{-2}] = -4L_0 + \frac{1}{2}C$, we obtain

$$\begin{aligned}-4(L_0 + c_0M_0) + \frac{1}{2}\text{ß}C &= \text{ß}\big([L_2, L_{-2}]\big) \\ &= -4u_2u_{-2}L_0 - 3(u_2v_{-2} + v_2u_{-2})Y_0 + \frac{1}{2}u_2u_{-2}C - 2u_2w_{-2}M_0 \\ &\quad -4v_2v_{-2}M_0 - 2w_2u_{-2}M_0.\end{aligned}$$

Comparing the coefficients of L_0, Y_0, using (3.30) and Lemma 3.1, one has (noting that $\sigma C \in C(\mathscr{L})$)

$$u_2 = u_{-2}^{-1} = u_1^2, \tag{3.31}$$

$$v_2 = -u_2 v_{-2} u_{-2}^{-1} = 2u_1 v_1, \tag{3.32}$$

$$ßC = C + 4(u_1 w_{-1} + 2v_1 v_{-1} + w_1 u_1^{-1} - u_2 w_{-2} - 2v_2 v_{-2} - w_2 u_{-2})M_0. \tag{3.33}$$

According to (3.28), (3.31) and (3.25), one obtains

$$u_n = u_1^n \quad \text{for any} \;\; n \in \mathbb{Z}^*. \tag{3.34}$$

Similarly, by (3.29), (3.32) and (3.26), one has

$$v_n = n u_1^{n-1} v_1 \quad \text{for any} \;\; n \in \mathbb{Z}^*. \tag{3.35}$$

Noting the equations (3.34) and (3.35), for $n \neq 0, \pm 1$, we can rewrite (3.27) as

$$(n-1)w_{n+1} = n u_1 w_n + (n-1)u_1^{n-1} v_1^2 - w_1 u_1^n. \tag{3.36}$$

For any $n \in \mathbb{Z}$, applying ß to both sides of $[L_1, Y_n] = (n - \frac{1}{2})Y_{n+1}$, one has

$$\begin{aligned}&(n-\frac{1}{2})(u_1 e'_n - e'_{n+1})Y_{n+1} + \big((n-1)v_1 e'_n + n u_1 e''_n - (n-\frac{1}{2})e''_{n+1}\big)M_{n+1}\\ &= (n-\frac{1}{2})\delta_{n+1,0}e_0 C.\end{aligned}$$

Comparing the coefficients of Y_{n+1}, M_{n+1} and C in the above equation, one has

$$e_0 = 0, \tag{3.37}$$

$$e'_{n+1} = u_1 e'_n, \tag{3.38}$$

$$(n-\frac{1}{2})e''_{n+1} = (n-1)v_1 e'_n + n u_1 e''_n. \tag{3.39}$$

Using induction on n in (3.38), one has

$$e'_n = u_1^n e'_0. \tag{3.40}$$

Applying ß to both sides of $[L_{-1}, Y_1] = \frac{3}{2}Y_0$, one has

$$\begin{aligned}\frac{3}{2}ß(Y_0) &= \frac{3}{2}e'_1 u_{-1} Y_0 + (2e'_1 v_{-1} + e''_1 u_{-1})M_0\\ &= \frac{3}{2}e'_1 u_1^{-1} Y_0 + (-2e'_1 u_1^{-2} v_1 + e''_1 u_1^{-1})M_0\\ &= \frac{3}{2}e'_0 Y_0 + (-2u_1 e'_0 u_1^{-2} v_1 + 2v_1 e'_0 u_1^{-1})M_0\\ &= \frac{3}{2}e'_0 Y_0.\end{aligned}$$

So we can rewrite $ßY_0$ as $ßY_0 = e'_0Y_0$, and $e''_0 = 0$. Using induction on n in (3.39), one has

$$e''_n = 2nu_1^{n-1}v_1e'_0. \tag{3.41}$$

By the equations (3.37), (3.40) and (3.41), for any $n \in \mathbb{Z}$, we can rewrite $ß(Y_n)$ as

$$ß(Y_n) = u_1^n e'_0 Y_n + 2nu_1^{n-1}v_1e'_0M_n. \tag{3.42}$$

For any $n \in \mathbb{Z}$, applying ß to both sides of $[L_1, M_n] = nM_{n+1}$, one has

$$nu_1f'_nM_{n+1} = nf'_{n+1}M_{n+1} + n\delta_{n+1,0}f_0C. \tag{3.43}$$

For $n = -1$, comparing the coefficients of M_0 and C in (3.43), one has

$$f'_{-1} = u_1^{-1}f'_0, \quad f_0 = 0. \tag{3.44}$$

For $n \neq 0, -1$, using (3.44) and comparing the coefficients of M_{n+1} and C in (3.43), one has

$$f'_{n+1} = u_1f'_n. \tag{3.45}$$

Applying ß to both sides of $[L_2, M_{-1}] = -M_1$ and comparing the coefficients of M_1, one has

$$f'_1 = u_2f'_{-1} = u_1^2u_1^{-1}f'_0 = u_1f'_0. \tag{3.46}$$

According to the equations (3.44), (3.46) and using induction on n in (3.45), one has

$$f'_n = u_1^nf'_0, \quad \forall\, n \in \mathbb{Z}. \tag{3.47}$$

By the equations (3.44) and (3.47), for any $n \in \mathbb{Z}$, we can rewrite $ß(M_n)$ as

$$ß(M_n) = u_1^nf'_0M_n. \tag{3.48}$$

Applying ß to $[Y_{-1}, Y_1] = 2M_0$ and comparing the coefficients of M_0, one has

$$f'_0 = {e'_0}^2. \tag{3.49}$$

By the equations (3.48) and (3.49), for any $n \in \mathbb{Z}$, we can rewrite $ß(M_n)$ as

$$ß(M_n) = u_1^n{e'_0}^2M_n. \tag{3.50}$$

By now, the results that we have obtained can be formulated as follows

$$\begin{aligned}
ß(L_n) &= \begin{cases} u_1^nL_n + nu_1^{n-1}v_1Y_n + w_nM_n, & n \neq 0; \\ L_0 + \frac{1}{2}(u_1w_{-1} - 2u_1^{-2}v_1^2 + u_1^{-1}w_1)M_0, & n = 0; \end{cases} && (3.51) \\
ß(Y_n) &= u_1^ne'_0Y_n + 2nu_1^{n-1}v_1e'_0M_n, \quad n \in \mathbb{Z}; && (3.52) \\
ß(M_n) &= u_1^n{e'_0}^2M_n, \quad n \in \mathbb{Z}; && (3.53) \\
ß(C) &= C + 4(u_1w_{-1} + 6u_1^{-2}v_1^2 + u_1^{-1}w_1 - u_1^2w_{-2} - u_1^{-2}w_2)M_0. && (3.54)
\end{aligned}$$

Replacing ß by $\psi_{e_0'^{-1}}\text{ß}_{u_1^{-1}}\text{ß}$ where $\text{ß}_{u_1^{-1}}$ and $\psi_{e_0'^{-1}}$ are defined in (3.2) and (3.4) and denoting ${e_0'}^{-1}u_1^{-1}v_1$ by v_1', ${e_0'}^{-2}u_1^{-n}w_n$ by w_n', we can rewrite the above equations as follows

$$\begin{aligned}
\text{ß}(L_n) &= \begin{cases} L_n + nv_1'Y_n + w_n'M_n, \quad n \neq 0; \\ L_0 + \frac{1}{2}(w_{-1}' - 2{v_1'}^2 + w_1')M_0, \quad n = 0; \end{cases} && (3.55) \\
\text{ß}(Y_n) &= Y_n + 2nv_1'M_n, \quad n \in \mathbb{Z}; && (3.56) \\
\text{ß}(M_n) &= M_n, \quad n \in \mathbb{Z}; && (3.57) \\
\text{ß}(C) &= C + 4(w_{-1}' + 6{v_1'}^2 + w_1' - w_{-2}' - w_2')M_0. && (3.58)
\end{aligned}$$

For $n \neq -1, 0$, applying σ to $[L_1, L_n] = (n-1)L_{n+1}$, we obtain

$$\begin{aligned}
\sigma\big([L_1, L_n]\big) &= (n-1)\big(L_{n+1} + (n+1)v_1'Y_{n+1} + w_{n+1}'M_{n+1}\big) \\
&= (n-1)L_{n+1} + \big(n(n-\frac{1}{2})v_1' + (\frac{n}{2}-1)v_1'\big)Y_{n+1} \\
&\quad + \big(nw_n' + n(n-1){v_1'}^2 - w_1'\big)M_{n+1}.
\end{aligned}$$

Comparing the coefficients of M_{n+1}, one has

$$(n-1)w_{n+1}' = nw_n' + n(n-1){v_1'}^2 - w_1'. \tag{3.59}$$

In particular, one has

$$w_{-2}' = w_{-1}' + 3{v_1'}^2 + \frac{w_{-1}' - w_1'}{2} = w_1' + 3{v_1'}^2 + \frac{-3(w_1' - w_{-1}')}{2}. \tag{3.60}$$

For $n \neq -1, 0$, applying σ to $[L_{-1}, L_{n+1}] = (n+2)L_n$, we obtain

$$\begin{aligned}
\sigma\big([L_{-1}, L_{n+1}]\big) &= (n+2)\big(L_n + nv_1'Y_n + w_n'M_n\big) \\
&= (n+2)L_n + \big((n+1)(n+\frac{3}{2})v_1' - \frac{n+3}{2}v_1'\big)Y_n \\
&\quad + \big((n+1)w_{n+1}' - (n+1)(n+2){v_1'}^2 + w_{-1}'\big)M_n.
\end{aligned}$$

Comparing the coefficients of M_n, one has

$$(n+1)w_{n+1}' = (n+2)w_n' + (n+1)(n+2){v_1'}^2 - w_{-1}'. \tag{3.61}$$

In particular, one has

$$w_2' = w_1' + 3{v_1'}^2 + \frac{w_1' - w_{-1}'}{2}. \tag{3.62}$$

For $n \neq -2, 0, \pm 1$, subtracting (3.59) from (3.61), one has

$$w_{n+1}' - w_n' = (2n+1){v_1'}^2 + \frac{w_1' - w_{-1}'}{2}. \tag{3.63}$$

According to the equations (3.60), (3.62) and (3.63), we obtain

$$w'_n = w'_1 + (n-1)(n+1){v'_1}^2 + \frac{(n-1)(w'_1 - w'_{-1})}{2}, \quad \forall\ n \in \mathbb{Z}^*. \tag{3.64}$$

If we denote v'_1, $\frac{w'_1 - w'_{-1}}{2}$, $w'_1 - {v'_1}^2 - \frac{w'_1 - w'_{-1}}{2}$ respectively by α, β, γ, then for any $n \in \mathbb{Z}$, the equations (3.55)–(3.58) can be rewritten as (3.19)–(3.22). The lemma follows. $\square$

From (3.19)—(3.22), one immediately see

$$\delta_{\alpha,\beta,\gamma}\delta_{\alpha',\beta',\gamma'} = \delta_{\alpha+\alpha',\beta+\beta',\gamma+\gamma'+2\alpha\alpha'}, \tag{3.65}$$

$$\delta_{\alpha,\beta,\gamma} = \delta_{\alpha',\beta',\gamma'} \iff \alpha = \alpha',\ \beta = \beta',\ \gamma = \gamma'. \tag{3.66}$$

For any

$$b = (\cdots, b_{-2}, b_{-1}, b_1, b_2, \cdots),\ c = (\cdots, c_{-2}, c_{-1}, c_1, c_2, \cdots) \in \mathbb{C}^\infty, \tag{3.67}$$

we denote

$$\xi_{b,c} = \exp\big(\sum_{j\neq 0} b_j \mathrm{ad} Y_j + \sum_{k\neq 0} c_k \mathrm{ad} M_k\big) \in \mathrm{Inn}(\mathscr{L}). \tag{3.68}$$

Then from the proofs of Lemmas 3.1–3.5, we see that every element $\sigma \in \mathrm{Aut}(\mathscr{L})$ can be uniquely expressed as $\big($the uniqueness can be seen from the fact that each data in $(b, c, i, u, w, \alpha, \beta, \gamma)$ is uniquely determined by $\sigma\big)$

$$\sigma = \xi_{b,c}\varepsilon^i \text{ß}_u \psi_w \delta_{\alpha,\beta,\gamma} \text{ for some } (b, c, i, u, w, \alpha, \beta, \gamma) \in \mathbb{C}^\infty \times \mathbb{C}^\infty \times \mathbb{Z}_2 \times \mathbb{C}^{*2} \times \mathbb{C}^3, \tag{3.69}$$

where $\mathbb{Z}_2 = \mathbb{Z}/2\mathbb{Z}$, and all notations can be found in (3.2)–(3.4), (3.19)—(3.22) and (3.68). Therefore there exists a one to one correspondence $f : \mathrm{Aut}(\mathscr{L}) \to \mathbb{C}^\infty \times \mathbb{C}^\infty \times \mathbb{Z}_2 \times \mathbb{C}^{*2} \times \mathbb{C}^3$,

$$f : \sigma \mapsto (b, c, i, u, w, \alpha, \beta, \gamma). \tag{3.70}$$

Suppose $\text{ß} = \xi_{b,c}\varepsilon^i \text{ß}_u \psi_w \delta_{\alpha,\beta,\gamma}$, $\text{ß}' = \xi_{b',c'}\varepsilon^{i'} \text{ß}_{u'} \psi_{w'} \delta_{\alpha',\beta',\gamma'} \in \mathrm{Aut}(\mathscr{L})$. Let

$$\text{ß}\text{ß}' = \text{ß}'' = \xi_{b'',c''}\varepsilon^{i''} \text{ß}_{u''} \psi_{w''} \delta_{\alpha'',\beta'',\gamma''}.$$

Then the data $(b'', c'', i'', u'', w'', \alpha'', \beta'', \gamma'') \in \mathbb{C}^\infty \times \mathbb{C}^\infty \times \mathbb{Z}_2 \times \mathbb{C}^{*2} \times \mathbb{C}^3$ can be determined by the following lemma.

Lemma 3.6. *Under the above notations, the following relations hold:*

$$w'' = ww', \tag{3.71}$$

$$i'' = i + i', \tag{3.72}$$

$$u'' = u^{(-1)^{i'}} u', \tag{3.73}$$

$$\gamma'' = w'^{-2}\gamma + \gamma', \tag{3.74}$$

$$\alpha'' = \frac{\alpha w'^{-1} + \alpha'}{2}, \tag{3.75}$$

$$\beta'' = w'^{-2}\beta + \alpha'^2 + \beta' + \gamma', \tag{3.76}$$

$$b''_j = b_j + (-1)^i w b'_{(-1)^i j} u^{(-1)^i j}, \tag{3.77}$$

$$\begin{aligned} c''_k = {} & c_k + w^2(-1)^i c'_{(-1)^i k} u^{(-1)^i k} + 2\alpha w^2 k b'_{(-1)^i k} u^{(-1)^i k} \\ & - \sum_{j\neq 0} \frac{(-1)^i w k^{-1}(k-j)(k-2j)\big(u^{(-1)^i j} b'_{(-1)^i j} b_{k-j} - u^{(-1)^i (k-j)} b_j b'_{(-1)^i (k-j)}\big)}{2}. \end{aligned} \tag{3.78}$$

Proof. Using the equation $\text{ßß}'(L_0) = \text{ß}''(L_0)$ and comparing the coefficients of L_0, Y_j $(j \neq 0)$, M_k $(k \neq 0)$ and M_0, we can deduce (3.72), (3.77), (3.78) and

$$w''^2\gamma'' = w^2\gamma + w^2 w'^2 \gamma'. \tag{3.79}$$

Applying both sides of $\text{ßß}' = \text{ß}''$ to Y_0 and comparing the coefficients of M_j $(j \neq 0)$, one can deduce (3.71). Then by (3.79), we obtain (3.74).

Similarly, applying both sides of $\text{ßß}' = \text{ß}''$ to M_1 and comparing the coefficients of $M_{(-1)^{i+i'}}$, one can deduce (3.73).

Using $\text{ßß}'(Y_1) = \text{ß}''(Y_1)$ and comparing the coefficients of $M_{(-1)^{i+i'}}$, one has (3.75).

Finally, applying both sides of $\text{ßß}' = \text{ß}''$ to L_1 and comparing the coefficients of $M_{(-1)^{i+i'}}$, one can deduce (3.76). ☐

Thus, we obtain the following theorem.

Theorem 3.7. *Under the map f defined in (3.70), the automorphism group* $\mathrm{Aut}(\mathscr{L})$ *is ismorphic to the group $\mathbb{C}^\infty \times \mathbb{C}^\infty \times \mathbb{Z}_2 \times \mathbb{C}^{*2} \times \mathbb{C}^3$ (where the elements in $\mathbb{C}^\infty$ is denoted as in (3.67)), whose group multiplication is given by*

$$(b, c, i, u, w, \alpha, \beta, \gamma) \cdot (b', c', i', u', w', \alpha', \beta', \gamma') = (b'', c'', i'', u'', w'', \alpha'', \beta'', \gamma''),$$

where the data $(b'', c'', i'', u'', w'', \alpha'', \beta'', \gamma'')$ is given by Lemma 3.6.

Corollary 3.8. $\mathrm{Inn}(\mathscr{L}) = \mathrm{Span}\big\{\exp\big(aL_0 + \sum_i b_i Y_i + \sum_{j\neq 0} c_j M_j\big) \mid a, b_i, c_j \in \mathbb{C}\big\}$.

Proof. The result follows immediately by noting from (3.1) that $\mathrm{Inn}(\mathscr{L})$ is generated by $\exp aL_0$, $\exp b_0 Y_0$ and $\xi_{b,c}$ (which is defined in (3.68)). ☐

References

[1] R. Farnsteiner, Derivations and central extensions of finitely generated graded Lie algebra, *J. Algebra*, **118** (1988), 33-45.

[2] C. Jiang, The holomorph and derivation algebra of infinite dimensional Heisenberg algebra, *Chinese J. Math.*, **17** (1997), 422-426.

[3] G. V. Kac, *Infinite Dimensional Lie Algebra*, 3rd ed, Cambrige, UK: Cambridge Univ. Press, (1990).

[4] D. Dokovic, K. Zhao, Derivations, isomorphisms and second cohomology of generalized Block algebras, *Alg. Colloq.*, **3** (1996), 245-272.

[5] M. Henkel, Schrödinger invariance and strongly anisotropic critical systems, *J. Stat. Phys.*, **75** (1994), 1023-1029.

[6] M. Henkel, Phenomenology of local scale invariance: from conformal invariance to dynamical scaling, *Nucl. Phys. B*, **641** (2002), 405-410.

[7] M. Henkel, J. Unterberger, Schrödinger invariance and space-time symmetries, *Nucl. Phys. B*, **660** (2003), 407-412.

[8] P. J. Hilton, U. Stammbach, *A Course in Homological Algebra.*, 2nd ed. New York: Springer-Verlag, (1997).

[9] V. Mazorchuk, K. Zhao, Classification of simple weight Virasoro modules with a finite-dimensional weight space, *J. Algebra*, **307** (2007), 209-214.

[10] C. Roger, J. Unterberger, The Schrödinger-Virasoro Lie group and algebra: from geometry to representation theory, preprint (arXiv:cond-mat/0601050), (2006).

[11] R. Shen, C. Jiang, The derivation algebra and the automorphism group of the twisted Heisenberg-Virasoro algebra, *Comm. Algebra*, **34** (2006), 2547-2558.

[12] Y. Su, Simple modules over the higher rank super-Virasoro algebras, *Lett. Math. Phys.*, **53** (2000), 263-272.

[13] J. Unterberger, On vertex algebra representations of the Schrödinger-Virasoro algebra, preprint (arXiv:cond-mat/0703214), (2007).

[14] L. Zhu, D. Meng, Some infinite dimensional complete Lie algebra, *China. Ann. Math.*, **21A**(3)(2000), 311-316.